

Combining Pharmacotherapy with Psychotherapeutic Management for the Treatment of Psychiatric Disorders

Ira D. Glick, M.D.
Stanford University School of Medicine

Richard Balon, M.D.
Wayne State University School of Medicine





Issue

What does psychotherapy add above what medication alone provides?:

- in what conditions?
- for which patients?
- at what phase?







Outline

- **Introduction**
 - Rationale and Model
 - Theoretical Outcomes
- Results
- Guidelines
- Summary and Clinical Implications
 -

Reasons Why Combined Therapy Not Delivered

- Many providers deliver one or the other
- Most insurers pay for non-integrated treatment
- Many professionals trained in one or the other
- Provider bias



Integrated vs. Collaborative Treatment

- **Integrated treatment** – pharmacotherapy and psychotherapy provided by one person
- **Collaborative (or split) treatment** – pharmacotherapy and psychotherapy provided by psychiatrist and therapist (psychologist, social worker, counselor)



Rationale for Combining Medication and Therapy

- Patients value psychotherapy
- Patient may not be on medication
- Etiology
 - Although biological, “stress” may precipitate episode
- Pathogenesis
 - Illness has effects on the family
- Treatment
 - To improve adherence
- It may work better than one modality (medication or therapy)

Reasons Why Psychotherapy is Important


- Some conditions have no effective pharmacotherapy treatment available
- Medication may be contraindicated
- Patient may not want to take medication (numerous reasons)
- Most patients have social and interpersonal problem accompanying Axis I disorders either as the source of or are the consequence of the illness

Outcomes of Combined Treatment


- Positive effects
- Negative effects
- No effect

Combined Treatment Outcome


No Additive Therapeutic Effect



Combined Treatment Outcome Positive Effect — Additive




Combined Treatment Outcome Positive Effect — Synergistic




Combined Treatment Outcome

Positive Effect — Facilitative



Medications Facilitate Accessibility to Psychotherapy





Outline

- Introduction
 - Rationale
 - Theoretical Outcomes
- **Results**
- Guidelines
- Summary and Clinical Implications

Results

- Controlled Research - examples TSS
- Guidelines and Algorithms
 - For schizophrenia - <50% received adequate treatment (Lehman et al, 1998)
 - For depression - consistent undertreatment (Keller et al, 1997)



Drug-Psychosocial Interactions in Schizophrenia

- Psychosocial treatments more effective when psychotic symptoms are controlled with drugs (May et al, 1968)
- Psychosocial treatments can be toxic when patients not adequately treated with drugs (Hogarty et al, 1974)
- Psychosocial treatments more effective when compliance is assured (Hogarty et al, 1979)

May PRA. *Treatment of Schizophrenia: A Comparative Study of Five Treatment Methods.*

New York, NY: Science House; 1968.

Hogarty GE et al. *Psychopharmacol Bull.* 1974;10:47.

Hogarty GE et al. *Arch Gen Psychiatry.* 1974;31:603-608.


Hogarty GE et al. *Arch Gen Psychiatry.* 1974;31:609-618.

Hogarty GE et al. *Arch Gen Psychiatry.* 1979;36:1283-1294.

Drug-Psychosocial Interactions in Schizophrenia

- Drugs may be more effective when compliance is enhanced by psychosocial treatment (Marder et al, 1996)
- Drugs and psychosocial treatments may affect different outcome domains, ie, drugs control symptoms and psychosocial treatments affect social adjustment (Marder et al, 1996)

Cumulative Relapse Rates in Schizophrenia



Trials Combining Medication With Psychosocial Treatments (Both Controlled)

	N	Medication (MED)	Psychosocial Treatment (PST)	Outcome	Result
Hogarty et al. (1973, 1974)	360	Chlorpromazine	Major role therapy	Relapse	>1 y: MED + ↑ PST
Hogarty et al. (1979)	105	Fluphenazine	Social therapy	Relapse	>1 y: MED + ↑ PST
Hogarty et al. (1986, 1991)	90	Fluphenazine	Family treatment, social skills training	Relapse, expressed emotion	1 y + 2 y: MED + ↑ PST
Schooler et al. (1997)	313	Fluphenazine	Psychoeducation vs family therapy	Rehospitalization symptoms	No difference between 2 PSTs
Marder et al. (1996)	80	Fluphenazine	Behavioral skills training, supportive group	Relapse, social adjustment	MED + PST adjustment ↑

Newer Antipsychotics and Quality of Life

- 1-year double-blind comparison of clozapine and haloperidol
- Clozapine-treated patients more likely to participate in psychosocial programs
- Participation in psychosocial treatment reduced symptoms and improved quality of life

***Table 1. Summary of Disorders In Which Psychotherapy Improves Outcome Over Medication Alone**

Disorder/ Syndrome	Treatments	Standard of Proof
Bipolar disorders	While pharmacological interventions are treatments of choice, psychosocial treatments, including psychoeducation, cognitive behavior therapy, IPSRT, and marital/family therapy, have shown the potential to increase medication adherence, improve quality of life, and enhance mechanisms for coping with stress in patients with bipolar disorder.	Several Type 2 and Type 3 studies of psychoeducation; three Type 1 studies of cognitive behavior therapy; one Type 1 study of IPSRT; and several Type 1 studies of marital/family therapy.
Childhood attention-deficit hyperactivity disorder (ADHD)	Combining intensive behavioral intervention with well-delivered pharmacological agents typically ranks better than either treatment component alone; this is the only modality that tends to normalize behavior patterns.	One very large-scale Type I clinical trial comparing behavioral intervention alone and medication alone with the two together
Major depressive disorder (MDD)	At least one major study lends strong support for the superior effectiveness of combined psychosocial and pharmacological treatment	One type 1 RCT



***Table 1. Summary of Disorders In Which Psychotherapy Improves Outcome Over Medication Alone Con't**

Disorder/ Syndrome	Treatments	Standard of Proof
Schizophrenia	Structured, educational family interventions help patients with schizophrenia maintain gains achieved with medication and customary case management	Over 20 Type 1 and Type 2 RCT's of educational family interventions
Post-traumatic stress disorder (PTSD)	Antidepressants reduced both PTSD symptoms and those of co-morbid conditions; they also made it easier for patients to benefit from psychotherapy, three varieties of antidepressants have been most commonly used	Several Type 1 and Type 2 RCT's
Sleep disorders	Behavioral interventions, including stimulus control, sleep restriction, relaxation strategies and cognitive behavioral therapy, have shown effectiveness, especially over the longer term in reducing sleep onset, decreasing awakenings, and increasing total sleep time; these behavioral interventions produce more sustained effects than pharmacological agents	A moderate number of Type 2 RCTs, in comparison to waitlist controls, partial behavioral interventions and pharmacological agents

*Studies are rated by quality of design, i. e. randomized controlled trials are the highest rated.

References are found in the text. From Nathan & Gorman, 2002, modified with permission



Table 1. Summary of Disorders in Which Psychotherapy Improves Outcome Over Medication Alone (cont' d)


Disorder/Syndrome	Treatments	Standard of Proof
Bulimia Nervosa (BN)		<ul style="list-style-type: none"> • A large number of Type 1 and Type 2 TCTs, utilizing placebo as comparison. • A very substantial number of Type 1 and Type 2 TCTs

Recent Studies Supporting Value of Adding Psychotherapy to Medication Alone for Adolescents

- Depression – TADS: CBT and Antidepressant
- Schizophrenia (early onset, ultra high risk) – McGorry: Psychoeducation and Antipsychotics
- Anxiety Disorder – CAMS: CBT and Antidepressant



Adjusted Predicted Mean Score
on Children's Depression Rating
Scale—Revised
(intention-to-treat analysis)^a



Chronic Depression

REVANT trial found no substantial short-term value of CBT or of supportive psychotherapy to results achieved by flexible, pharmacotherapy alone



Combination Therapy: Treatment Strategies

- For partial responders, unhelpful to ask whether to use drug or psychotherapy
- Comb Rx should be encouraged in those NOT responding to either modality (alone)



Pharmaco Rx > Psycho Rx

- Schizophrenia
- Dysthymia



Adding Pharmacologic Rx to Psycho Rx > Psycho Rx (Alone)

- MDD
- Social Phobia
- Bulimia



Adding Psycho Rx to Pharmaco Rx > Drug Rx Alone

- Depressive Disorder
- Schizophrenia
- Panic Disorder
- Bulimia

Leucht et al, 2014





Outline

- Introduction
 - Rationale
 - Theoretical Outcomes
- Results
- **Guidelines**
- Summary and Clinical Implications

Presumed Mechanism of Action

- “Correct the presumed biochemical deficit, but then use the window of opportunity provided by suppression of the symptom to remold both cognitions and behavior.”

Malcolm Lader, 1996

- “I believe that biologically mediated impairments in abstract thinking, “executive” problem solving, and mood reactivity (i.e., the ability to lift one’s spirits in response to something savory or encouraging will be found to be the clinical culprits that result from the biological alterations described above.”

Michael Thase, 1998



The Concept of Long-term Management

- **The aim:** to decrease symptoms and improve quality of life.
- **Process:** interactive among patient, family/sign other and MD.
- **Method:** in person and by phone p r n, but on-going regular contact



General Guidelines for Combined Therapy: I –Diagnosis

- Make a DSM diagnosis
- Make a family systems diagnosis
- Make an individual dynamic diagnosis
- Formulate case



General Guidelines for Combined Therapy: II – Goals

- Select appropriate modalities and their combination
- Develop specific goals for each modality
- Decide about sequencing
- Be aware of, and enquire about, side effects of each modality as well as their interactive effects



Hierarchy of Treatment Goals in Medical Psychotherapy of Schizophrenia

- **Acute Phase**

- Medical/neuropsychiatric assessment
 - Rapid symptom reduction
 - Reduce impact of episode on friends, family, housing, activities

- **Convalescent Phase**

- Gain trust/alliance with family/caregivers
 - Assess and mobilize social support
 - Ensure human service needs are met (food, clothing housing)
 - Ensure safety and predictability of environment



Hierarchy of Treatment Goals in Medical Psychotherapy of Schizophrenia (cont' d)

- **Adaptive Plateau**

Establish therapeutic alliance/supportive treatment routine

Achieve effective maintenance medication regime

- **Stable Plateau**

Psychoeducation: Promote illness self- management strategies, awareness of relationship between stress and symptoms

Rehabilitation: Teach adaptive competencies



Hierarchy of Treatment Goals in Medical Psychotherapy of Schizophrenia (cont' d)

- **Selected Patients Near Recovery**
 - Encourage introspection, self and other awareness
 - Encourage improved interpersonal relations, productivity, conflict management, self-understanding, self-concept



General Guidelines for Combined Therapy: III – Sequencing of Combined Treatments

- Step 1 - Establish an alliance
- Step 2 - For psychotic disorders, start medication early
- Step 3 - Add individual intervention as patient is able to participate
- Step 4 - Add family intervention early. Start with psychoeducation and referral to appropriate group depending on DSM diagnosis



General Guidelines for Combined Therapy: III – Sequencing of Combined Treatments (cont)

- Step 5 - Add family dynamic and systemic intervention as patient stabilizes
- Step 6 - Rehabilitation in maintenance phase
- Step 7 - Do not add another modality, if first intervention (vs second) adequate for efficacy.



Guidelines for Structure of Session

- Assumption: Minimum 15 minutes, maximum 60 minutes
- mean 30 minutes
- Divide session in three parts:
- Part I - 5 minutes:
 - ask global questions
 - then ask about side effects and target symptoms
(compared to baseline)
 - do psychoeducation
 - adjust medication



Guidelines for Structure of Session (cont)

- Part 2 - 20 minutes:
 - explore life events
 - explore issues of transference to “pill” and to psychotherapy
 - explore issues of countertransference



Guidelines for Structure of Session (cont)

- Part 3 - 5 minutes, integration of:

- patient issues
- family issues
- provide final prescription of medication and summarize therapy, discussed in psychotherapy, review possible side effects, complications



Guidelines for Medications

- **New Medications:**
- **New Strategies:**
 1. **Adequate (rather than inadequate dose), but minimal effective dose (MED)**
 2. **Continuous (rather than targeted)**
 3. **Standard (moderate) dose (rather than low dose)**
 4. **I-M long-acting agents for non-compliant patients (rather than po)**



Medications (cont)

- **Pharmacotherapeutic Alliance (PA, Gutheil):**
 1. **Defined as the manner in which active efforts are made by the physician to enlist, recruit, involve patient in a collaboration around prescribing meds**
 2. **Characteristics - flexible, prescriptive stance**
 3. **Objective - establishment and maintenance of the PA**
 4. **Process - shared enquiry, shared goals, mutual participation in both the experience and observing the process of using medication**



Guidelines for Psychotherapy

- Which type in which phase
 - family intervention early if patient cognitively impaired
 - individual intervention as patient able to participate
 - combine both with family session as needed in maintenance phases

Guidelines for Family Intervention

- 1) A positive approach and genuine working relationship between the therapist and family.
- 2) The provision of family therapy in a table, structured format with the availability of additional contacts with therapists if necessary.
- 3) A focus on improving stress and coping in the “here and how,” rather than dwelling on the past

Guidelines for Family Intervention (cont)

- 4) Encouragement of respect for interpersonal boundaries within the family
- 5) The provision of information about the biological nature of schizophrenia in order to reduce blaming of the patient and family guilt
- 6) Use of behavioral techniques, such as breaking down goals into manageable steps.
- 7) Improving communication among family members

Summary of Effects of Family Intervention and Pharmacological Intervention

Including Rehabilitation

- Family Intervention
 - Education
 - Communication Skills
 - Problem Solving Skills
 - Resolution of Dynamic and Systems Issues
- Pharmacological Intervention
 - Normalize Illness
 - Suppress Symptoms



Guidelines for Effective Individual Intervention in Psychotic Disorder

- First - make an alliance with patient and family
- Help patient maintain self-esteem regardless of illness
- Focus on improving adherence
- Distinguish among and manage among:
 - Objective psychopathology
 - Psychodynamic issues
 - Personality conflicts/deficits from patient's life history



Problems Emanating from Misuse of the Psychodynamic Model in Axis I Disorders

- 1) Emphasizing the patient's personality conflicts and character pathology over the patient's overt symptom manifestations.
- 2) Misdiagnosing objective dependency as psychodynamic dependency.
- 3) Misdiagnosis of lifelong disability as psychodynamic psychopathology.
- 4) Labeling cognitive and other changes resulting from Axis-I disorders as a personality disorder, usually “narcissistic, borderline or histrionic.”
- 5) Treating with psychoanalytic psychotherapy with the expectation that Axis-I symptoms will subside after personality conflicts are resolved.
- 6) Overvaluing the effectiveness of therapist role, i.e. transference over the natural history of disease or medication noncompliance



Psychoeducation for Combined Treatment

- Provide systematic information (repetitively) to both patient and to family about
 - signs and symptoms
 - diagnosis
 - treatment - both medication and psychotherapy
 - prognosis with and without treatment
- Aim for behavioral change in both patient and family - don't just provide information



Rehabilitation

- When
 - Post-acute episode (Harding — over 2-3 decades function may increase)
- How much
 - As tolerated
- What
 - Program for Assertive Community Outreach (PACT)
 - Case management
 - Social skills training (Lieberman)
 - Sheltered workshops
 - Vocational rehabilitation
- Who
 - Positive and negative symptom issues
- Why
 - To achieve options that patients desire

Combined Therapy:

Advantages

- 1) For those patients biologically-oriented, psychotherapy promotes a sense of increased collaboration and targets intrapsychic and interpersonal problems
- 2) For those patients psychologically-oriented, medication response relieves hopelessness associated with lack of improvement in psychotherapy as well as targeting primary S_x of illness
- 3) Faster response than either modality alone
- 4) Family therapy can ↑ medication compliance
- 5) Individual therapy can ↑ medication compliance
- 6) Medication can ↑ psychotherapy compliance



Combined Therapy: Disadvantages


- 1) With medication, ↑ risk for side effects and early termination of all therapies
- 2) With psychotherapy, perceived need for medication decreased (“I can solve this on my own”)
- 3) Splitting





Outline

- Introduction
 - Rationale
 - Theoretical Outcomes
- Results
- Guidelines
- **Summary and Clinical Implications**
 -




Nature via Nurture


A revolution is sweeping the field of biology that holds that the influences of nature and nurture are so inextricably linked that it is difficult to speak of them as distinct forces that shape who we are. We now know that our environment can change us only if we are genetically predisposed to change and that our genes are powerless if they are not primed by the environment. When it comes to understanding our fate, we can no longer study the effect of genes of the environment without considering the interaction between them.

C.T. Gross, 2004





The New Yorker 1992




MANKOFF

"Oh, yeah? Well, I think you're the one with the biochemical imbalance."



Quality Treatment Equation



Reference

- Glick ID: Adding psychotherapy to pharmacotherapy: Data, benefits, and guidelines for integration. *Am J Psychotherapy*. 2004 58:186-208



